

543
NPSOR-91-02

NAVAL POSTGRADUATE SCHOOL

Monterey, California



CLASSIFICATION OF INTERMITTENT DEPENDENT OBSERVATIONS

P. A. Jacobs
and
D. P. Gaver

October 1990

Approved for public release; distribution is unlimited.

Prepared for:
Naval Ocean Systems Center
San Diego, CA 92152

FedDocs
D 208.14/2
NPS-OR-91-02

2nd Doc
202 14/2
1PS-OP-21-02

**NAVAL POSTGRADUATE SCHOOL,
MONTEREY, CALIFORNIA**

Rear Admiral R. W. West, Jr.
Superintendent

Harrison Shull
Provost

This report was prepared in conjunction with research funded by the Naval Ocean Systems Center, San Diego, California.

Unclassified

Security Classification of this page

DUDLEY KNOX LIBRARY
NAVAL POSTGRADUATE SCHOOL
MONTEREY, CALIFORNIA 93943-5002DUDLEY KNOX LIBRARY
NAVAL POSTGRADUATE SCHOOL
MONTEREY CA 93943-5101

REPORT DOCUMENTATION PAGE

1a Report Security Classification UNCLASSIFIED		1b Restrictive Markings	
2a Security Classification Authority		3 Distribution Availability of Report	
2b Declassification/Downgrading Schedule		Approved for public release; distribution is unlimited	
4 Performing Organization Report Number(s) NPSOR-91-02		5 Monitoring Organization Report Number(s)	
6a Name of Performing Organization Naval Postgraduate School	6b Office Symbol (If Applicable) OR	7a Name of Monitoring Organization Naval Ocean Systems Center	
6c Address (city, state, and ZIP code) Monterey, CA 93943-5000		7b Address (city, state, and ZIP code) San Diego, CA 92152	
8a Name of Funding/Sponsoring Organization Naval Ocean Systems Center	8b Office Symbol (If Applicable)	9 Procurement Instrument Identification Number N6600190WR00093	
8c Address (city, state, and ZIP code) San Diego, CA 92152		10 Source of Funding Numbers	
		Program Element Number	Project No
		Task No	Work Unit Accession No
11 Title (Include Security Classification) Classification of Intermittent Dependent Observations			
12 Personal Author(s) P. A. Jacobs and D. P. Gaver			
13a Type of Report Technical	13b Time Covered From To	14 Date of Report (year, month, day) 1990, October	15 Page Count 38
16 Supplementary Notation The views expressed in this paper are those of the author and do not reflect the official policy or position of the Department of Defense or the U.S. Government.			
17 Cosati Codes		18 Subject Terms (continue on reverse if necessary and identify by block number)	
Field	Group	Subgroup	
		Bayesian paradigm, Gaussian autoregressive process; Cauchy autoregressive process; classification; model misspecification; robustness	
19 Abstract (continue on reverse if necessary and identify by block number)			
<p>Each of J items has a characteristic Signature which varies in time. At time 0, the value of a Signature and the identity of the corresponding item are known. No further values of Signatures are observed until a later time $t > 0$. At time t a Signature associated with an unknown item is observed. The problem is to estimate the identity of the item whose Signature is observed at time t. The estimation procedure studied is to estimate the identity of the item that is associated with the Signature at time t to be that one which maximizes the posterior probability of being associated with the observed Signature. Univariate and multivariate Gaussian and univariate Cauchy autoregressive processes are considered as models for the Signatures. The robustness of the univariate Gaussian, (respectively Cauchy), procedure when applied to Cauchy, (respectively Gaussian) data is studied. The results suggest that the Gaussian classification procedure is biased towards classifying the Signature observed at time t as being associated with the <i>same</i> item that is associated with the Signature at time 0. The Cauchy procedure is biased towards classifying a Signature observed at time t as being associated with a <i>different</i> item than the one associated with the Signature at time 0.</p>			
20 Distribution/Availability of Abstract <input checked="" type="checkbox"/> unclassified/unlimited <input type="checkbox"/> same as report <input type="checkbox"/> DTIC users		21 Abstract Security Classification Unclassified	
22a Name of Responsible Individual D. P. Gaver		22b Telephone (Include Area code) (408) 646-2605	22c Office Symbol OR/Gv, OR/Jc

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted

security classification of this page

All other editions are obsolete

Unclassified

DUDLEY KNOX LIBRARY
NAVAL POSTGRADUATE SCHOOL
MONTEREY, CALIFORNIA 93943-5002

CLASSIFICATION OF INTERMITTENT DEPENDENT OBSERVATIONS

by
P. A. Jacobs and D. P. Gaver

1. THE PROBLEM

Consider the following classification problem. Suppose there are J items (e.g., diseases) each of which has a characteristic Signature which varies in time; the Signature of Item i is

$$Y_i(t) = \theta_i + X_i(t) \quad i = 1, 2, \dots, J \quad t = 0, 1, 2, \dots \quad (1.1)$$

For the moment $\{X_i(t)\}$ is an unspecified multivariate (or univariate) stochastic process, but one that stays near θ_i in finite time and has some stationary or steady-state behavior. In many cases, paths of $X_i(t)$ will appear somewhat “continuous,” so successive $X_i(t)$ ’s are not well-modeled as iid random variables. One could think of $Y_i(t)$ as physical indices characteristic of a particular disease, e.g., blood pressure, heart-beat pattern, cholesterol levels. Examples from equipment reliability are also of interest; here physical indices might be vibration, variations in heat level, oil leakage, and even fuel consumption in the case of engines.

In many circumstances $Y_i(t)$ is only observable occasionally, at times unrelated to the value of $Y_i(t)$ but driven by other forces such as the scheduling of a routine physical exam or system inspection. Suppose that the Signature and the identity of the item associated with the Signature are both observed at time $t = 0$, on such an occasion. Suppose that, later on, however, only the Signature of an item is observed. The first question is: What is the probability that, given the Signature value observed, its originating item is any particular one of the J candidates?

In Gaver and Jacobs [1989], the processes $\{X_i(t)\}$ are assumed to be univariate Gaussian and a Bayesian classification procedure is studied. In this

paper, Section 2 assumes $\{X_i(t)\}$ are multivariate normal autoregressive processes. In Section 3, $\{X_i(t)\}$ is a univariate Cauchy autoregressive process whose marginal distribution has longer tails than the Gaussian. A Bayesian classification procedure for the Cauchy data is studied. In Section 4, we study the behavior of the univariate Cauchy and Gaussian classification procedures when autoregressive data having the wrong marginal distribution are presented to them. The results suggest that the Gaussian classification procedure is biased towards classifying a Signature produced at time t as being associated with the *same* item that produced the Signature at time 0. The Cauchy classification procedure is biased towards classifying a Signature produced at time t as being associated with a *different* item than the one producing the Signature at time 0. These effects are strongest for small times t . The largest number of misclassifications occur for small times t when the Gaussian classification procedure is presented with Cauchy data and a *different item* is associated with the Signature at time t than the item associated with the Signature at time 0; in this situation the Gaussian procedure is relatively less sensitive to the change in the item associated with the Signature. Misclassifications by the Cauchy classification procedure are modest in comparison to this extreme case.

In summary, it is important to realize that the performance of a Bayesian classification procedure can be influenced by its underlying distributional assumptions. A classification procedure based on Gaussian distributional assumptions can be reluctant to classify a new observation coming from a different item as being associated with a new item. A classification procedure based on Cauchy distributional assumptions can be reluctant to classify a new observation which comes from the same item as that being associated with the same item. Hence, if there is uncertainty about the underlying distribution of the data, it might be better to combine results of several classification procedures based on different distributional assumptions.

2. THE MULTIVARIATE NORMAL CASE

2.1 The Classification Question

Assume for illustration that the Signature of Item j is multivariate AR(1):

$$\mathbf{Y}_j(t) = \boldsymbol{\theta}_j + \mathbf{X}_j(t) \quad (2.1)$$

where $\boldsymbol{\theta}_j$, $\mathbf{Y}_j(t)$, and $\mathbf{X}_j(t)$ are d -dimensional column vectors. The process $\{\mathbf{X}_j(t)\}$ is a vector AR(1) process

$$\mathbf{X}_j(t) = \mathbf{A}_j \mathbf{X}_j(t-1) + \mathbf{E}_j(t) \quad (2.2)$$

where \mathbf{A}_j is a $d \times d$ matrix and $\{\mathbf{E}_j(t)\}$ is a sequence of d -dimensional column vectors which are independent multivariate normal with mean $\mathbf{0}$ and variance-covariance matrix $\boldsymbol{\Lambda}_j$. The variance-covariance matrix for $\mathbf{X}_j(t+1)$ is

$$\boldsymbol{\Gamma}_j(t+1) = E[\mathbf{X}(t+1)\mathbf{X}^T(t+1)] = \mathbf{A}_j \boldsymbol{\Gamma}_j(t) \mathbf{A}_j^T + \boldsymbol{\Lambda}_j. \quad (2.3)$$

We will assume \mathbf{A}_j and $\boldsymbol{\Lambda}_j$ are such that there is a finite unique solution to the equation

$$\boldsymbol{\Gamma}_j = \mathbf{A}_j \boldsymbol{\Gamma}_j \mathbf{A}_j^T + \boldsymbol{\Lambda}_j. \quad (2.4)$$

Assume $\mathbf{X}_j(0)$ has a normal distribution with mean $\mathbf{0}$ and variance-covariance matrix $\boldsymbol{\Gamma}_j$. It follows that $\{\mathbf{X}_j(t)\}$ is a stationary sequence with mean $\mathbf{0}$ and variance-covariance matrix $\boldsymbol{\Gamma}_j$.

The conditional distribution of $\mathbf{X}_j(t)$ given $\mathbf{X}_j(0) = \mathbf{x}$ is multivariate normal with mean $\mathbf{A}_j^t \mathbf{x}$ and variance-covariance matrix

$$\Lambda_j(t) = \sum_{n=0}^{t-1} \mathbf{A}_j^n \Lambda_j \left(\mathbf{A}_j^n \right)^T. \quad (2.5)$$

Thus, $\Gamma_j = \lim_{t \rightarrow \infty} \Lambda_j(t)$.

The conditional distribution of the actually observable $\mathbf{Y}_j(t)$ given $\mathbf{Y}_j(0)=\mathbf{y}(0)$ is multivariate normal with mean $\boldsymbol{\theta}_j + \mathbf{A}_j^t(\mathbf{y}(0) - \boldsymbol{\theta}_j)$ and variance-covariance matrix $\Lambda_j(t)$.

Operational Scenario: There are, potentially, J items. Let $C(t)$ be the identity of the item whose Signature is observed at time t . Put $p_j(t) \equiv P\{C(t)=j\}$. Assume that it is known that the Signature observed at time 0 comes from Item i ; that is, $C(0) = i$ and $\mathbf{Y}(0) = \mathbf{Y}_i(0) = \mathbf{y}(0)$. If it has been a long time since a Signature from item i has been observed, it is reasonable to suppose that

$$\begin{aligned} P\{C(0) = i, \mathbf{Y}(0) = \mathbf{y}(0)\} \\ = p_i(0) \left((2\pi)^d |\Gamma_i| \right)^{-0.5} \exp \left\{ -\frac{1}{2} (\mathbf{y}(0) - \boldsymbol{\theta}_i)^T \Gamma_i^{-1} (\mathbf{y}(0) - \boldsymbol{\theta}_i) \right\} \end{aligned} \quad (2.6)$$

the long-run or steady-state distribution. Further,

$$\begin{aligned} P\{C(t) = i, \mathbf{Y}(t) = \mathbf{y}(t) | C(0) = i, \mathbf{Y}(0) = \mathbf{y}(0)\} \\ = p_i(t) \left[(2\pi)^d |\Sigma_i(t)| \right]^{-0.5} \exp \left\{ -\frac{1}{2} (\mathbf{y}(t) - \mathbf{m}_i(t))^T \Sigma_i(t)^{-1} (\mathbf{y}(t) - \mathbf{m}_i(t)) \right\} \end{aligned} \quad (2.7)$$

where

$$\mathbf{m}_i(t) = \boldsymbol{\theta}_i + \mathbf{A}_i^t (\mathbf{y}(0) - \boldsymbol{\theta}_i) \quad (2.8)$$

and

$$\Sigma_i(t) = \Lambda_i(t). \quad (2.9)$$

For $j \neq i$, we will assume the conditional distribution of $\mathbf{Y}(t)$ given $C(0) = i$, $\mathbf{Y}(0) = \mathbf{y}(0)$, $C(t) = j$ is multivariate normal with mean $\mathbf{m}_j(t) = \boldsymbol{\theta}_j$ and variance-covariance matrix $\boldsymbol{\Gamma}_j \equiv \boldsymbol{\Sigma}_j(t)$, since it is still a long time since a Signature from Item j is observed.

It now follows that

$$\begin{aligned} P\{\mathbf{Y}(t) \in d\mathbf{y}(t) | C(0) = i, \mathbf{Y}(0) = \mathbf{y}(0)\} \\ = \sum_{j=1}^J p_j(t) \left[(2\pi)^d |\boldsymbol{\Sigma}_j(t)| \right]^{-0.5} \exp \left\{ -\frac{1}{2} (\mathbf{y}(t) - \mathbf{m}_j(t))^T \boldsymbol{\Sigma}_j(t)^{-1} (\mathbf{y}(t) - \mathbf{m}_j(t)) \right\}. \end{aligned} \quad (2.10)$$

Thus, the posterior probability of the identity of the item associated with the Signature is

$$\begin{aligned} P\{C(t) = j | C(0) = i, \mathbf{Y}(0) = \mathbf{y}(0), \mathbf{Y}(t) = \mathbf{y}(t)\} = \\ p_j(t) |\boldsymbol{\Sigma}_j(t)|^{-0.5} \exp \left\{ -\frac{1}{2} (\mathbf{y}(t) - \mathbf{m}_j(t))^T \boldsymbol{\Sigma}_j(t)^{-1} (\mathbf{y}(t) - \mathbf{m}_j(t)) \right\} \\ \times \left[\sum_{k=1}^J p_k(t) |\boldsymbol{\Sigma}_k(t)|^{-0.5} \exp \left\{ -\frac{1}{2} (\mathbf{y}(t) - \mathbf{m}_k(t))^T \boldsymbol{\Sigma}_k(t)^{-1} (\mathbf{y}(t) - \mathbf{m}_k(t)) \right\} \right]^{-1}. \end{aligned} \quad (2.11)$$

2.2 The Probability of an Incorrect Classification

In this section we assume that the item that is associated with the Signature at time t given the last complete observation at time 0 will be estimated to be that one which maximizes the posterior probability (2.11).

For a simple illustration we will suppose that there are only $J=2$ possible items with known parameters $\boldsymbol{\theta}_1$ and $\boldsymbol{\theta}_2$.

Given $\mathbf{Y}(0)=\mathbf{y}(0)$, $C(0)=1$, and $C(t) = 1$, the conditional distribution of $\mathbf{Y}(t)$ is multivariate normal with mean

$$\mathbf{m}_1(t) = \boldsymbol{\theta}_1 + \mathbf{A}_1^t (\mathbf{y}(0) - \boldsymbol{\theta}_1) \quad (2.12)$$

and variance-covariance matrix

$$\boldsymbol{\Sigma}_1(t) = \sum_{k=0}^{t-1} \left(\mathbf{A}_1^k \right)^T \boldsymbol{\Lambda}_1 \mathbf{A}_1^k. \quad (2.13)$$

Let the matrices $\mathbf{H}_i(t)$ and \mathbf{H}_i be such that

$$\mathbf{H}_i(t) \mathbf{H}_i(t)^T = \boldsymbol{\Sigma}_i(t) \quad (2.14)$$

and

$$\mathbf{H}_i \mathbf{H}_i^T = \boldsymbol{\Gamma}_i. \quad (2.15)$$

It follows that

$$\mathbf{Y}(t) \stackrel{d}{=} \mathbf{m}_1(t) + \mathbf{H}_1(t) \mathbf{U} \quad (2.16)$$

where \mathbf{U} is a d -dimensional column vector each of whose components are independent standard normal random variables; the notation $\stackrel{d}{=}$ means equal in distribution. Thus, given $\mathbf{Y}(0) = \mathbf{y}(0)$, $C(0) = 1$, $C(t) = 1$,

$$(\mathbf{Y}(t) - \mathbf{m}_1(t))^T \boldsymbol{\Sigma}_1(t)^{-1} (\mathbf{Y}(t) - \mathbf{m}_1(t)) \stackrel{d}{=} \mathbf{U}^T \mathbf{U} \quad (2.17)$$

and

$$\begin{aligned} & (\mathbf{Y}(t) - \mathbf{m}_2(t))^T \boldsymbol{\Sigma}_2(t)^{-1} (\mathbf{Y}(t) - \mathbf{m}_2(t)) \\ & \stackrel{d}{=} (\mathbf{m}_1(t) - \mathbf{m}_2(t) + \mathbf{H}_1(t) \mathbf{U})^T \boldsymbol{\Sigma}_2(t)^{-1} (\mathbf{m}_1(t) - \mathbf{m}_2(t) + \mathbf{H}_1(t) \mathbf{U}) \end{aligned} \quad (2.18)$$

where $\mathbf{m}_2(t) = \boldsymbol{\theta}_2$ and $\boldsymbol{\Sigma}_2(t) = \boldsymbol{\Gamma}_2$. Thus, the probability of a misclassification is

$$\begin{aligned}
& P\{\text{classify the item as } 2 | C(0) = 1, Y(0) = y(0), C(t) = 1\} \\
& = P\left\{p_2(t)|\Sigma_2(t)|^{-0.5} \exp\left\{-\frac{1}{2}(\mathbf{m}_1(t) - \mathbf{m}_2(t) + \mathbf{H}_1(t)\mathbf{U})^T \Sigma_2(t)^{-1}(\mathbf{m}_1(t) - \mathbf{m}_2(t) + \mathbf{H}_1(t)\mathbf{U})\right\}\right. \\
& \quad \left. > p_1(t)|\Sigma_1(t)|^{-0.5} \exp\left\{-\frac{1}{2}\mathbf{U}^T \mathbf{U}\right\}\right\} \\
& = P\left\{\frac{p_2(t)}{p_1(t)}\left(\frac{|\Lambda_1(t)|}{|\Gamma_2|}\right)\right\}^{\frac{1}{2}} > \exp\left\{-\frac{1}{2}\mathbf{U}^T \mathbf{U} + \frac{1}{2}(\mathbf{a}(t) + \mathbf{H}_1(t)\mathbf{U})^T \Gamma_2^{-1}(\mathbf{a}(t) + \mathbf{H}_1(t)\mathbf{U})\right\} \quad (2.19)
\end{aligned}$$

where

$$\mathbf{a}(t) = \boldsymbol{\theta}_1 + \mathbf{A}_1^t(\mathbf{y}(0) - \boldsymbol{\theta}_1) - \boldsymbol{\theta}_2. \quad (2.20)$$

Example: Assume $\Lambda_i = \Lambda, \mathbf{A}_i = \mathbf{A}$ for $i = 1, 2$, and $p_1(t) = p_2(t)$; then

$$\begin{aligned}
& P\{\text{wrong classification} | C(0) = 1, Y(0) = y(0), C(t) = 1\} \\
& = P\left\{\left(\frac{|\Lambda(t)|}{|\Gamma|}\right)^{\frac{1}{2}} > \exp\left\{-\frac{1}{2}\mathbf{U}^T \mathbf{U} + \frac{1}{2}(\mathbf{a}(t) + \mathbf{H}(t)\mathbf{U})^T \Gamma^{-1}(\mathbf{a}(t) + \mathbf{H}(t)\mathbf{U})\right\}\right\} \quad (2.21)
\end{aligned}$$

where

$$\Lambda(t) = \sum_{k=0}^{t-1} \mathbf{A}^k \Lambda (\mathbf{A}^k)^T \equiv \mathbf{H}(t)\mathbf{H}(t)^T;$$

$\Gamma = \lim_{t \rightarrow \infty} \Lambda(t) \equiv \mathbf{H}\mathbf{H}^T$ is the solution to the equation

$$\Gamma = \mathbf{A}\Gamma\mathbf{A}^T + \Lambda;$$

and

$$\mathbf{a}(t) = \boldsymbol{\theta}_1 + \mathbf{A}^t(\mathbf{y}(0) - \boldsymbol{\theta}_1) - \boldsymbol{\theta}_2.$$

Note that as $t \rightarrow \infty$

$$\begin{aligned}
& P\{\text{wrong classification} \mid C(0) = 1, Y(0) = y(0), C(t) = 1\} \\
& \rightarrow P\left\{1 > \exp\left\{-\frac{1}{2}U^T U + \frac{1}{2}(\theta_1 - \theta_2 + HU)^T \Gamma^{-1}(\theta_1 - \theta_2 + HU)\right\}\right\} \\
& = P\left\{1 > \exp\left\{\frac{1}{2}(\theta_1 - \theta_2)^T (\mathbf{H}\mathbf{H}^T)^{-1}(\theta_1 - \theta_2) + (\theta_1 - \theta_2)^T (\mathbf{H}^T)^{-1}U\right\}\right\} \\
& = P\left\{-\frac{1}{2}(\theta_1 - \theta_2)^T (\mathbf{H}\mathbf{H}^T)^{-1}(\theta_1 - \theta_2) > (\theta_1 - \theta_2)^T (\mathbf{H}^T)^{-1}U\right\}. \quad (2.22)
\end{aligned}$$

A Simulation

Table 1 gives the results of a simulation experiment for the case $\theta_1 = (1,1)^T$

$$A = \begin{bmatrix} 0.1 & 0.9 \\ 0.4 & 0.5 \end{bmatrix}$$

and

$$\Lambda = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

In this case,

$$\Gamma = \begin{bmatrix} 5.4 & 3.8 \\ 3.8 & 4.5 \end{bmatrix}.$$

Figure 1 shows contours from a bivariate normal distribution having mean θ_1 and variance-covariance matrix Γ .

In each replication two independent vector random variables are generated; one is $Y(0)$ which has a normal distribution with mean θ_1 and variance-covariance matrix Γ ; the other is U , whose components are two independent standard normal random variables. For each time $t = 1, 2, \dots, 40$, $Y(t)$ is calculated as

$$Y(t) = m(t) + H(t)U. \quad (2.23)$$

with $m(t) = \theta_1 + A^t (y(0) - \theta_1)$; $Y(t)$ has the same distribution as a Signature from Item 1 when the Signature at time 0 is also from Item 1. There are 1000 replications. Table 1 presents the fraction of replications for which the incorrect classification is made of Item 2 being the one producing the Signature at time t ; that is those replications for which

$$\left(\frac{|\Lambda(t)|}{|\Gamma|} \right)^{\frac{1}{2}} > \exp \left\{ \frac{1}{2} (y(t) - \theta_2)^T \Gamma^{-1} (y(t) - \theta_2) - \frac{1}{2} (y(t) - \theta_1)^T \Lambda(t)^{-1} (y(t) - \theta_1) \right\}. \quad (2.24)$$

Note that the fractions are not independent since common random numbers are used.

The contours of the distribution in Figure 1 suggest that it is more likely to make a misclassification if $\theta_2 = (2,2)^T$ than if $\theta_2 = (-2,2)^T$; the fractions in Table A support this. The fractions in Table A also suggest that the probability of misclassification is an increasing function of t . This observation is supported by the fact that the variances of the components of $Y(t)$ increase as t increases.

TABLE A. FRACTION OF MISCLASSIFICATION

$$\theta_1 = (1,1)^T$$

Time:	1	2	3	4	5	10	20	30	40
$\theta_2 = (2,2)^T$	0.10	0.13	0.16	0.20	0.21	0.30	0.39	0.41	0.41
$\theta_2 = (-2,2)^T$	0.04	0.06	0.06	0.07	0.07	0.09	0.09	0.09	0.09

3. CAUCHY UNIVARIATE MODEL

In this section we consider Bayesian classification for a time series model having marginal distributions with a longer tail than the Gaussian distribution.

We assume that

$$Y_i(t) = \theta_i + X_i(t)$$

with

$$X_i(t) = \rho_i X_i(t-1) + \varepsilon_i(t)$$

where $|\rho_i| < 1$; $\{\varepsilon_i(t)\}$ are independent sequences of independent identically distributed Cauchy random variables with location parameter 0 and precisions $\left[(1-|\rho_i|)\alpha_i\right]^{-0.5}$; and $X_i(0)$ has a Cauchy distribution with parameters 0 and $\alpha_i^{-0.5}$. Under these assumptions $\{X_i(t); t = 0, 1, 2, \dots\}$ is a stationary sequence of random variables with marginal Cauchy distribution having parameters 0 and $\alpha_i^{-0.5}$.

It follows that

$$\begin{aligned} & P\{Y_i(0) \in dy(0), Y_i(t) \in dy_i(t)\} \\ &= \frac{1}{\pi} \alpha_i \left[\alpha_i^2 + (y(0) - \theta_i)^2 \right]^{-1} \frac{1}{\pi} \alpha_i (1 - |\rho_i|^t) \left[\left(\alpha_i (1 - |\rho_i|^t) \right)^2 + \left(y(t) - \theta_i - \rho_i^t (y(0) - \theta_i) \right)^2 \right]^{-1}. \end{aligned} \quad (3.1)$$

Let $C(t)$ denote the identity of the item associated with the Signature at time t and put $P\{C(t)=i\}=p_i(t)$; then

$$\begin{aligned} & P\{Y(t) \in dy(t), C(t) = i | C(0) = i, Y(0) = y(0)\} \\ &= p_i(t) \frac{1}{\pi} \alpha_i (1 - |\rho_i|^t) \left[\left(\alpha_i (1 - |\rho_i|^t) \right)^2 + \left(y(t) - \theta_i - \rho_i^t (y(0) - \theta_i) \right)^2 \right]^{-1} \end{aligned}$$

$$\equiv p_i(t) \frac{1}{\pi} \alpha_i(t) \left[\alpha_i(t)^2 + (y(t) - m_i(t))^2 \right]^{-1}. \quad (3.2)$$

Thus,

$$P\{Y(t) \in dy(t) | C(0) = i, Y(0) = y(0)\} = \sum_{j=1}^J p_j(t) \frac{1}{\pi} \alpha_j(t) \left[\alpha_j(t)^2 + (y(t) - m_j(t))^2 \right]^{-1}$$

where $\alpha_i(t)$ and $m_i(t)$ are defined in (3.2) and it is natural to define $m_j(t) = \theta_j$ and $\alpha_j(t) = \alpha_j$ for $j \neq i$. Hence, given item i is associated with the Signature at time 0, the posterior probability that item j is associated with the Signature observed at time t is

$$\begin{aligned} P\{C(t) = j | C(0) = i, Y(0) = y(0), Y(t) = y(t)\} \\ = p_j(t) \alpha_j(t) \left[\alpha_j(t)^2 + (y(t) - m_j(t))^2 \right]^{-1} \\ \times \left\{ \sum_{k=1}^J p_k(t) \alpha_k(t) \left[\alpha_k(t)^2 + (y(t) - m_k(t))^2 \right]^{-1} \right\}^{-1}. \end{aligned} \quad (3.3)$$

3.2 The Probability of Making an Incorrect Classification

In this section we assume that the item associated with the Signature at time t given the last complete observation at time 0 is estimated to be that one which maximizes the posterior probability (3.3). For simplicity we will suppose there are $J=2$ possible items with known parameters θ_1 and θ_2 .

First

$$\begin{aligned} P\{Y(t) \in dy(t) | Y(0) = y(0), C(0) = 1, C(t) = j\} \\ = \frac{1}{\pi} \alpha_j(t) \left[\left(\alpha_j(t) \right)^2 + (y(t) - m_j(t))^2 \right]^{-1} \end{aligned} \quad (3.4)$$

where

$$\alpha_1(t) = \alpha_1 \left[1 - |\rho_1|^t \right]; \alpha_2(t) = \alpha_2 \quad (3.5)$$

$$m_1(t) = \theta_1 + \rho_1^t (y(0) - \theta_1); m_2(t) = \theta_2. \quad (3.6)$$

Note that given $Y(0) = y(0)$, $C(0) = 1$ and $C(t) = 1$,

$$Y(t) \stackrel{d}{=} \left[\theta_1 + \rho_1^t (y(0) - \theta_1) \right] + \left(1 - |\rho_1|^t \right) \alpha_1 W \equiv m_1(t) + \alpha_1(t) W$$

where W is a Cauchy random variable with location parameter 0 and precision 1. Hence, the probability of making the incorrect classification of estimating Item 2 as being associated with the Signature at time t given Item 1 is responsible for Signatures at time 0 and time t and $Y(0) = y(0)$ is

$$\begin{aligned} & P\{\text{Classify as Item 2} | C(0) = 1, Y(0) = y(0), C(t) = 1\} \\ &= P\left\{ \frac{p_2(t)}{p_1(t)} \alpha_2(t) \left[\alpha_2(t)^2 + (Y(t) - \theta_2)^2 \right]^{-1} > \alpha_1(t) \left[\alpha_1(t)^2 + (Y(t) - m_1(t))^2 \right]^{-1} \mid C(0) = 1, Y(0) = y(0), C(t) = 1 \right\} \\ &= P\left\{ \frac{p_2(t)}{p_1(t)} \frac{\alpha_2(t)}{\alpha_1(t)} \left[\alpha_2^2 + (m_1(t) + \alpha_1(t)W - \theta_2)^2 \right]^{-1} > \left[\alpha_1(t)^2 + (\alpha_1(t)W)^2 \right]^{-1} \mid C(0) = 1, Y(0) = y(0), C(t) = 1 \right\} \\ &= P\left\{ \frac{p_2(t)}{p_1(t)} \alpha_2(t) \alpha_1(t) > \frac{\left[\alpha_2^2 + (m_1(t) + \alpha_1(t)W - \theta_2)^2 \right]}{1 + W^2} \mid C(0) = 1, Y(0) = y(0), C(t) = 1 \right\}. \end{aligned} \quad (3.7)$$

Note that as $t \rightarrow 0$, $\alpha_1(t) \rightarrow 0$, and $m_1(t) \rightarrow y(0)$. Hence, the conditional probability of a wrong classification tends to

$$P\left\{ 0 > \frac{\left[\alpha_2^2 + (y(0) - \theta_2)^2 \right]}{1 + W^2} \right\} = 0. \quad (3.8)$$

As $t \rightarrow \infty$, $\alpha_1(t) \rightarrow \alpha_1$, $m_1(t) \rightarrow \theta_1$ and the conditional probability of a wrong classification tends to

$$P\left\{\frac{p_2(\infty)}{p_1(\infty)}\alpha_2\alpha_1(1+W^2) > \left[\alpha_2^2 + (\alpha_1 W + \theta_1 - \theta_2)^2\right]\right\}. \quad (3.9)$$

If $\alpha_2 = \alpha_1 = \alpha$ and $p_2(\infty) = p_1(\infty)$, then as $t \rightarrow \infty$

$$\begin{aligned} &P\{\text{incorrect classification} | Y(0) = y(0), C(0) = 1, C(t) = 1\} \\ &= P\left\{\alpha^2(1+W^2) > \alpha^2\left(1 + \left[W + \frac{(\theta_1 - \theta_2)}{\alpha}\right]^2\right)\right\} \\ &= P\left\{W^2 > \left[W + \frac{(\theta_1 - \theta_2)}{\alpha}\right]^2\right\} \\ &= P\left\{W > \left|\frac{\theta_1 - \theta_2}{2\alpha}\right|\right\} \end{aligned} \quad (3.10)$$

which increases as α increases and decreases as $|\theta_1 - \theta_2|$ increases.

4. ARE BAYESIAN CLASSIFICATION PROCEDURES ROBUST?

In this section the robustness of the univariate Cauchy and Gaussian classification procedures against misspecification of the form of the marginal distribution will be studied.

4.1 Gaussian Data.

In this subsection we assume that the Signatures of the Items form Gaussian time series. In particular we assume that

$$Y_i(t) = \theta_i + X_i(t) \quad (4.1)$$

with

$$X_i(t+1) = \rho_i X_i(t) + \varepsilon_i(t) \quad (4.2)$$

where $\{\varepsilon_i(t)\}$ are independent identically distributed normal random variables with mean 0 and variance σ_i^2 and $|\rho_i| < 1$. The independent random variable $X_i(0)$ has a normal distribution with mean 0 and variance $\sigma_i(\infty)^2 = \sigma_i^2 / (1 - \rho_i^2)$. Thus $\{X_i(t), t \geq 0\}$ is a stationary sequence of normal random variables with mean 0 and variance $\sigma_i(\infty)^2$. Let $C(t)$ be the identity of the Item associated with the Signature at time t .

As was shown in Gaver and Jacobs (1989), the conditional distribution of $Y(t)$ given $Y(0)=y(0)$, $C(0)=i$, $C(t)=i$ is normal with mean

$$m_i(t) = \theta_i + (y(0) - \theta_i)\rho_i^t \quad (4.3)$$

and standard deviation

$$\sigma_i(t) = \sigma_i(\infty)\sqrt{1 - \rho_i^{2t}}. \quad (4.4)$$

For simplicity we will assume $P\{C(t) = i\} = p_i(t) \equiv p(t)$ and there are 2 Items with parameters θ_1 and θ_2 ; thus, $p(t) = \frac{1}{2}$

Suppose the Cauchy procedure is used to estimate the identity of the Item associated with the Signature at time t ; that is, the Item which maximizes the posterior probability (3.3) is the estimate of the Item associated with the Signature. Hence, the probability of an incorrect classification is

$$\begin{aligned}
 & P\{\text{Classify as Item 2} | C(0) = 1, Y(0) = y(0), C(t) = 1\} \\
 &= P\left\{ \frac{\alpha_2(t)}{\alpha_1(t)} \left[\alpha_2(t)^2 (m_1(t) + \sigma_1(t)Z - \theta_2)^2 \right]^{-1} > \left[\alpha_1(t)^2 + (\sigma_1(t)Z)^2 \right]^{-1} \right\} \\
 &= P\left\{ \alpha_2(t)\alpha_1(t) \left[\alpha_2(t)^2 + (m_1(t) + \sigma_1(t)Z - \theta_2)^2 \right]^{-1} > \left[1 + \frac{(\sigma_1(t)Z)^2}{\alpha_1(t)} \right]^{-1} \right\} \\
 &= P\left\{ \alpha_2\alpha_1(1-|\rho_1|^t) \left[\alpha_2(t)^2 + (m_1(t) + \sigma_1(t)Z - \theta_2)^2 \right]^{-1} > \left[1 + \left(\frac{\sigma_1(\infty)\sqrt{1-\rho_1^{2t}}}{\alpha_1(1-|\rho_1|^t)} Z \right)^2 \right]^{-1} \right\} \\
 & \tag{4.5}
 \end{aligned}$$

where Z is a standard normal random variable.

Note that as $t \rightarrow 0$

$$\begin{aligned}
 & P\{\text{Classify as Item 2} | C(0) = 1, Y(0) = y(0), C(t) = 1\} \\
 &= P\left\{ \alpha_2\alpha_1(1-|\rho_1|^t) \left[\alpha_2^2 + (m_1(t) + \sigma_1(t)Z - \theta_2)^2 \right]^{-1} > \left[1 + \left[\frac{\sigma_1(\infty)}{\alpha_1} \right]^2 \frac{1-\rho_1^{2t}}{(1-|\rho_1|^t)^2} Z^2 \right]^{-1} \right\}
 \end{aligned}$$

$$\begin{aligned}
&= P \left\{ \alpha_2 \alpha_1 (1 - |\rho_1|^t) \left[\alpha_2^2 + (m_1(t) + \sigma_1(t)Z - \theta_2)^2 \right]^{-1} > \left[1 + \left[\frac{\sigma_1(\infty)}{\alpha_1} \right]^2 \frac{1 + |\rho_1|^t}{1 - |\rho_1|^t} Z^2 \right]^{-1} \right\} \\
&= P \left\{ \alpha_1 \alpha_2 (1 - |\rho_1|^t) \left[\alpha_2^2 + (m_1(t) + \sigma_1(t)Z - \theta_2)^2 \right]^{-1} > (1 - |\rho_1|^t) \left[(1 - |\rho_1|^t) + \left[\frac{\sigma_1(\infty)}{\alpha_1} \right]^2 (1 + |\rho_1|^t) Z^2 \right]^{-1} \right\} \\
&= P \left\{ \left[\alpha_2^2 + (m_1(t) + \sigma_1(t)Z - \theta_2)^2 \right] < \alpha_1 \alpha_2 \left[(1 - |\rho_1|^t) + \left[\frac{\sigma_1(\infty)}{\alpha_1} \right]^2 (1 + |\rho_1|^t) Z^2 \right] \right\} \\
&\rightarrow P \left\{ \frac{1}{\alpha_1 \alpha_2} \left[\alpha_2^2 + (y(0) - \theta_2)^2 \right] < \left[\frac{\sigma_1(\infty)}{\alpha_1} \right]^2 2Z^2 \right\} \\
&= P \left\{ \frac{\alpha_1}{2\alpha_2 \sigma_1(\infty)^2} \left[\alpha_2^2 + (y(0) - \theta_2)^2 \right] < Z^2 \right\}. \tag{4.6}
\end{aligned}$$

Thus, the conditional probability of an incorrect classification does not tend to 0 as $t \rightarrow \infty$ as it would if the correct model were used; see Gaver and Jacobs (1989) (3.6).

Note that as $t \rightarrow \infty$

$$\begin{aligned}
&P\{\text{Classify as Item 2} | C(0) = 1, Y(0) = y(0), C(t) = 1\} \\
&= P \left\{ \alpha_2 \alpha_1 \left[\alpha_2^2 + (\theta_1 + \sigma_1(\infty)Z - \theta_2)^2 \right]^{-1} > \left[1 + \left(\frac{\sigma_1(\infty)}{\alpha_1} Z \right)^2 \right]^{-1} \right\}. \tag{4.7}
\end{aligned}$$

If $\alpha_1 = \alpha_2 = 1$, then the above equals

$$P \left\{ \left[1 + (\sigma_1(\infty)Z + \theta_1 - \theta_2)^2 \right]^{-1} > \left[1 + (\sigma_1(\infty)Z)^2 \right]^{-1} \right\}$$

$$\begin{aligned}
&= P\left\{(\sigma_1(\infty)Z)^2 > (\sigma_1(\infty)Z + \theta_1 - \theta_2)^2\right\} \\
&= P\left\{Z^2 > \left(Z + \frac{(\theta_1 - \theta_2)}{\sigma_1(\infty)}\right)^2\right\} \\
&= P\left\{Z > \frac{|\theta_1 - \theta_2|}{2\sigma_1(\infty)}\right\} \tag{4.8}
\end{aligned}$$

which is the same as if the correct model had been used to make the decision; see (3.9) of Gaver and Jacobs (1989).

Now we consider the case in which a different Item is associated with the Signature at time t than the one associated with the Signature at time 0. Once again for simplicity we assume $\sigma_1 = \sigma_2 = \sigma$, $\rho_1 = \rho_2 = \rho$ with $|\rho| < 1$ and for the Cauchy model $\alpha_1 = \alpha_2 = \alpha$. Let $\sigma(\infty) = \sigma / \sqrt{1-\rho^2}$ and $\sigma(t) = \sigma(\infty)\sqrt{1-\rho^{2t}}$. We will assume Item 1 is associated with the Signature at time 0 and Item 2 is associated with the Signature at time t .

For the Gaussian classification procedure of Gaver and Jacobs (1989), the probability of an incorrect classification is

$$\begin{aligned}
&P\{\text{classify as 1} \mid Y(0) = y(0), C(0) = 1, C(t) = 2\} \\
&= P\{(1-\rho^{2t})^{-1/2} \exp\{-\frac{1}{2} [\theta_2 + \sigma(\infty)Z - (\theta_1 + (y(0)-\theta_1)\rho^t)]^2 / \sigma(\infty)^2 (1-\rho^{2t})\} \\
&\quad > \exp\{-\frac{1}{2} (\theta_2 + \sigma(\infty)Z - \theta_2)^2 / \sigma(\infty)^2\}\} \\
&= P\{-\frac{1}{2} \ln (1-\rho^{2t}) - \frac{1}{2} [\theta_2 + \sigma(\infty)Z - (\theta_1 + (y(0)-\theta_1)\rho^t)]^2 / \{\sigma(\infty)^2 (1-\rho^{2t})\} \\
&\quad > -\frac{1}{2} (\sigma(\infty)Z)^2 / \sigma(\infty)^2\} \\
&= P\{(1-\rho^{2t})\sigma(\infty)^2 \ln (1-\rho^{2t}) + [\theta_2 + \sigma(\infty)Z - (\theta_1 + (y(0)-\theta_1)\rho^t)]^2 \\
&\quad < (1-\rho^{2t}) (\sigma(\infty)Z)^2\} \tag{4.9}
\end{aligned}$$

where Z is a standard normal random variable. As $t \rightarrow 0$

$$P\{\text{classify as 1} \mid Y(0) = y(0), C(0) = 1, C(t) = 2\} \rightarrow P\left\{\left(Z + \frac{\theta_2 - y(0)}{\sigma(\infty)}\right)^2 < 0\right\} = 0; \quad (4.10)$$

that is, if the Item associated with the Signature at time t is different than the one associated with the Signature at time 0, then as $t \rightarrow 0$, the probability of an incorrect classification using the Gaussian procedure on Gaussian data tends to 0.

As $t \rightarrow \infty$, the probability of an incorrect classification,

$$\begin{aligned} P\{\text{classify as 1} \mid Y(0) = y(0), C(0) = 1, C(t) = 2\} \\ \rightarrow P\left\{\left(Z + \frac{\theta_2 - \theta_1}{\sigma(\infty)}\right)^2 < Z^2\right\} = P\left\{Z < \frac{-|\theta_2 - \theta_1|}{2\sigma(\infty)}\right\}. \end{aligned} \quad (4.11)$$

Suppose now the Cauchy classification procedure is used on the Gaussian data with Item 2 associated with the Signature at time t and Item 1 associated with the Signature at time 0. The probability of an incorrect classification

$$\begin{aligned} P\{\text{classify as 1} \mid Y(0) = y(0), C(0) = 1, C(t) = 2\} \\ = P\{\alpha(1 - |\rho|^t) [\alpha(1 - |\rho|^t)]^2 + [\theta_2 + \sigma(\infty)Z - [\theta_1 + (y(0) - \theta_1)\rho^t]]^2\}^{-1} > \alpha[\alpha^2 + (\theta_2 + \sigma(\infty)Z - \theta_2)^2]^{-1} \} \\ = P\{(1 - |\rho|^t) [\alpha^2 + (\sigma(\infty)Z)^2] > [(\alpha(1 - |\rho|^t))^2 + [\theta_2 + \sigma(\infty)Z - [\theta_1 + (y(0) - \theta_1)\rho^t]]^2]\}. \end{aligned} \quad (4.12)$$

As $t \rightarrow 0$

$$\begin{aligned} P\{\text{classify as 1} \mid Y(0) = y(0), C(0) = 1, C(t) = 2\} \\ \rightarrow P\{0 > [\theta_2 + \sigma(\infty)Z - y(0)]^2\} = 0; \end{aligned} \quad (4.13)$$

that is, as $t \rightarrow 0$, the probability of a correct classification for the Cauchy procedure tends to 1 for the case in which the Item associated with the Signature at time t is different from the one associated with the Signature at time 0, even though the data are Gaussian.

As $t \rightarrow \infty$

$P(\text{classify as 1} \mid Y(0) = y(0), C(0) = 1, C(t) = 2)$

$$\rightarrow P\left\{Z^2 > \left(Z + \frac{\theta_2 - \theta_1}{\sigma(\infty)}\right)^2\right\} = P\left\{Z < -\frac{|\theta_2 - \theta_1|}{2\sigma(\infty)}\right\}. \quad (4.14)$$

Hence, as $t \rightarrow \infty$ the probability of an incorrect identification tends to the same normal tail probability for both the Cauchy and Gaussian classification procedures.

Thus, for the two limiting cases $t \rightarrow 0$ and $t \rightarrow \infty$, both the Cauchy and Gaussian procedures have the same misclassification probabilities for the scenario in which the Item associated with the Signature at time t is different than the one associated with the Signature at time 0. Note that these are theoretical limiting results with all parameters known.

To investigate further the behavior of the two classification procedures on Gaussian data when Item 1 is associated with the Signature at time 0 and Item 2 is associated with the Signature at time t , let

$$g_t(y(0), Z) = (\theta_2 + \sigma(\infty)Z - [\theta_1 + (y(0) - \theta_1)\rho^t])^2.$$

The conditional probability of an incorrect classification by the Gaussian procedure is from (4.9)

$$P\{\text{Classify as Item 1} | Y(0) = y(0), C(0) = 1, C(t) = 2\}$$

$$\begin{aligned} &= P\{(1-\rho^{2t})\sigma(\infty)^2 \ln(1-\rho^{2t}) < (1-\rho^{2t}) (\sigma(\infty)Z)^2 - g_t(y(0), Z)\} \\ &\geq P\{(1-\rho^{2t})\sigma(\infty)^2 \ln(1-\rho^{2t}) < (1-|\rho|^t) (\sigma(\infty)Z)^2 - g_t(y(0), Z)\} \\ &\geq P\{-\alpha^2(1-|\rho|^t) | \rho|^t < (1-|\rho|^t) (\sigma(\infty)Z)^2 - g_t(y(0), Z)\} \end{aligned}$$

for t sufficiently close to 0. From (4.12) it follows that the conditional probability of misclassification for the Cauchy procedure is

$$P\{(1-|\rho|^t)(\sigma(\infty)Z)^2 - g_t(y(0), Z) > \alpha^2(1-|\rho|^t) [1-|\rho|^{t-1}]\}.$$

Hence for t sufficiently small, the incorrect Cauchy procedure will tend to have fewer misclassifications than the Gaussian procedure applied to Gaussian data in the scenario in which different Items are producing the Signatures at time 0 and t .

4.2 Cauchy Data

In this subsection we assume the Signatures form time series with Cauchy marginal distributions as in Section 3. In particular, we assume that

$$Y_i(t) = \theta_i + X_i(t) \tag{4.15}$$

with

$$X_i(t+1) = \rho_i X_i(t) + \varepsilon_i(t) \tag{4.16}$$

where $\{\varepsilon_i(t)\}$ are independent identically distributed Cauchy random variables with location 0 and precision $[(1-|\rho_i|)\alpha_i]^{-0.5}$ with $|\rho_i| < 1$. The independent random variable $X_i(0)$ has a Cauchy distribution with location 0 and precision $\alpha_i^{-1/2}$. Under these assumptions $\{X_i(t)\}$ is a stationary sequence of random variables with marginal Cauchy distribution having parameters 0 and $\alpha_i^{-1/2}$.

Further, the conditional distribution of $Y(t)$ given $Y(0) = y(0)$, $C(0) = i$, $C(t) = i$ is Cauchy with location parameter

$$m_i(t) = \theta_i + \rho_i^t(y(0) - \theta_i) \quad (4.17)$$

and precision parameter $\alpha_i(t)^{-1/2}$ with

$$\alpha_i(t) = \alpha_i(1 - |\rho_i|^t). \quad (4.18)$$

Let $C(t)$ be the identity of the Item associated with the Signature at time t .

For simplicity we will assume there are two items with parameters θ_1 and θ_2 . Further $P\{C(t) = i\} = p_i(t) \equiv p(t)$.

Suppose the Gaussian procedure of Gaver and Jacobs (1989) is used to estimate the identity of the item associated with the Signature at time t ; that is, the item which maximizes (2.12) of Gaver and Jacobs (1989) is the estimate of the Item associated with the Signature at time t . Hence, the probability of an incorrect classification is

$$\begin{aligned} & P\{\text{Classify as Item 2} | C(0) = 1, Y(0) = y(0), C(t) = 1\} \\ &= P\left\{ \frac{1}{\sigma_2(\infty)} \exp\left\{ -\frac{1}{2} \left[\frac{Y(t) - \theta_2}{\sigma_2(\infty)} \right]^2 \right\} > \frac{1}{\sigma_1(t)} \exp\left\{ -\frac{1}{2} \left[\frac{Y(t) - m_1(t)}{\sigma_1(t)} \right]^2 \right\} \middle| \begin{array}{l} C(0) = 1, \\ Y(0) = y(0), \\ C(t) = 1 \end{array} \right\} \quad (4.19) \end{aligned}$$

where

$$m_1(t) = \theta_1 + \rho_1^t(y(0) - \theta_1) \quad (4.20)$$

$$\sigma_1(t) = \sigma_1(\infty) \sqrt{1 - \rho_1^{2t}} \quad (4.21)$$

$$\sigma_1(\infty) = \sigma_1 / \sqrt{1 - \rho_1^2}; \sigma_2(\infty) = \sigma_2 / \sqrt{1 - \rho_2^2} \quad (4.22)$$

and σ_i , $i = 1, 2$ are the assumed standard deviations of the normal distributions. We will assume $\sigma_1 = \sigma_2 = \sigma$; $\rho_1 = \rho_2 = \rho$; $\alpha_1 = \alpha_2$. Hence,

$$P\{\text{Classify as Item 2} | C(0) = 1, Y(0) = y(0), C(t) = 1\} \\ = P\left\{\frac{1}{\sigma(\infty)} \exp\left\{-\frac{1}{2}\left(\frac{m_1(t) + \alpha(t)W - \theta_2}{\sigma(\infty)}\right)^2\right\} > \frac{1}{\sigma(t)} \exp\left\{-\frac{1}{2}\left(\frac{\alpha(t)W}{\sigma(t)}\right)^2\right\}\right\} \quad (4.23)$$

where W is a Cauchy random variable with location parameter 0 and precision 1.

$$P\{\text{Classify as Item 2} | C(0) = 1, Y(0) = y(0), C(t) = 1\} \\ = P\left\{\sqrt{1 - \rho^{2t}} > \exp\left\{-\frac{1}{2} \frac{(\alpha[1 - |\rho|^t])^2}{(1 - \rho^{2t})\sigma(\infty)^2} W^2 + \frac{1}{2} \frac{(\rho^t(y(0) - \theta_1) + (\theta_1 - \theta_2) + \alpha(t)W)^2}{\sigma(\infty)^2}\right\}\right\} \\ = P\left\{\sqrt{1 - \rho^{2t}} > \exp\left\{-\frac{1}{2} \frac{\alpha^2[1 - |\rho|^t]}{(1 + |\rho|^t)\sigma(\infty)^2} W^2 + \frac{1}{2} \frac{(\rho^t(y(0) - \theta_1) + (\theta_1 - \theta_2) + \alpha(t)W)^2}{\sigma(\infty)^2}\right\}\right\} \quad (4.24)$$

$$\xrightarrow{t \rightarrow 0} P\left\{0 > \exp\left\{\frac{1}{2} \frac{(y(0) - \theta_2)^2}{\sigma(\infty)^2}\right\}\right\} = 0;$$

thus, the probability of misclassification tends to zero as $t \rightarrow 0$ even though the incorrect model is being used; the correct Cauchy procedure also has a probability of misclassification tending to zero as $t \rightarrow 0$. As $t \rightarrow \infty$

$$P\{\text{Classify as Item 2} | C(0) = 1, Y(0) = y(0), C(t) = 1\} \\ = P\left\{1 > \exp\left\{-\frac{1}{2} \frac{\alpha^2 W^2}{\sigma(\infty)^2} + \frac{1}{2} \frac{[\theta_1 - \theta_2 + \alpha W]^2}{\sigma(\infty)^2}\right\}\right\}$$

$$\begin{aligned}
&= P\left\{1 > \exp\left\{\frac{1}{2} \frac{1}{\sigma(\infty)^2} \left[(\theta_1 - \theta_2)^2 + 2(\theta_1 - \theta_2)\alpha W\right]\right\}\right\} \\
&P\left\{0 > \frac{1}{2}(\theta_1 - \theta_2)^2 + (\theta_1 + \theta_2)\alpha W\right\} \\
&= P\left\{W < -\frac{1}{2} \frac{1}{\alpha} |\theta_1 - \theta_2|\right\}
\end{aligned}$$

which is the same as (3.10) the corresponding probability when the correct Cauchy procedure is used.

To further explore the behavior as $t \rightarrow 0$, let

$$g_t(W, y(0)) = [\rho^t(y(0) - \theta_1) + (\theta_1 - \theta_2) + \alpha(1 - |\rho|^t)W]^2$$

and

$$B(t) = \alpha^2(1 - |\rho|^t)W^2.$$

For t small (4.24) becomes

$$\begin{aligned}
&P\{\text{Classify as Item 2} | Y(0) = y(0), C(0) = 1, C(t) = 1\} \\
&= P\{\sigma(\infty)^2(1 + |\rho|^t) \ln(1 - \rho^{2t}) + B(t) > (1 + |\rho|^t) g_t(W, y(0))\} \\
&\leq P\{-\alpha^2 |\rho|^t + B(t) > (1 + |\rho|^t) g_t(W, y(0))\} \\
&\leq P\{-\alpha^2 |\rho|^t + B(t) > g_t(W, y(0))\} \\
&= P\{\alpha^2(1 - |\rho|^t) + B(t) > \alpha^2 + g_t(W, y(0))\} \\
&= P\{(\alpha(1 - |\rho|^t))^2 + (1 - |\rho|^t) B(t) > (1 - |\rho|^t) [\alpha^2 + g_t(W, y(0))]\} \\
&= P\{[\alpha^2 + g_t(W, y(0))]^{-1} > \alpha(1 - |\rho|^t) [(\alpha(1 - |\rho|^t))^2 + (\alpha(1 - |\rho|^t)W)^2]^{-1}\}
\end{aligned}$$

which is the conditional probability of misclassification for the Cauchy procedure on Cauchy data. Hence for small t , the incorrect Gaussian procedure will tend to have fewer misclassifications than the correct Cauchy procedure for the scenario in which the same item is associated with the Signature at both times.

Now we consider the case in which the Item associated with the Signature at time t is different than the one associated with the Signature at time 0. Once again for simplicity we assume $\alpha_1 = \alpha_2 = \alpha$, $\rho_1 = \rho_2 = \rho$ with $|\rho| < 1$ and for the Gaussian model $\sigma_1 = \sigma_2 = \sigma$. Let $\sigma(\infty) = \sigma / (1 - \rho^2)^{0.5}$ and $\sigma(t) = \sigma(\infty) (1 - \rho^{2t})^{0.5}$. We will assume Item 1 is associated with the Signature at time 0 and Item 2 is associated with the Signature at time t .

For the Gaussian classification procedure of Gaver and Jacobs (1989) the probability of an incorrect classification is

$$\begin{aligned}
 & P\{\text{classify as 1} \mid Y(0) = y(0), C(0) = 1, C(t) = 2\} \\
 &= P\{(1 - \rho^{2t})^{-0.5} \exp\{-\frac{1}{2} [\theta_2 + \alpha W - (\theta_1 + \rho^t(y(0) - \theta_1))]^2 / \sigma(\infty)^2 (1 - \rho^{2t})\} \\
 &\quad > \exp\{-\frac{1}{2} (\theta_2 + \alpha W - \theta_2)^2 / \sigma(\infty)^2\}\} \\
 &= P\{-\frac{1}{2} \ln(1 - \rho^{2t}) - \frac{1}{2} [\theta_2 + \alpha W - (\theta_1 + \rho^t(y(0) - \theta_1))]^2 / \{\sigma(\infty)^2 (1 - \rho^{2t})\} \\
 &\quad > -\frac{1}{2} (\alpha W)^2 / \sigma(\infty)^2\} \\
 &= P\{\sigma(\infty)^2 (1 - \rho^{2t}) \ln(1 - \rho^{2t}) + (\theta_2 + \alpha W - (\theta_1 + \rho^t(y(0) - \theta_1)))^2 \\
 &\quad < (\alpha W)^2 (1 - \rho^{2t})\} \tag{4.26}
 \end{aligned}$$

where W is a standard Cauchy random variable.

As $t \rightarrow 0$, the probability of an incorrect classification,

$$P\{\text{classify as 1} \mid Y(0) = y(0), C(0) = 1, C(t) = 2\}$$

$$\rightarrow P\{(\theta_2 + \alpha W - y(0))^2 < 0\} = 0.$$

Hence as $t \rightarrow 0$, the probability of an incorrect classification tends to 0 for the Gaussian procedure on Cauchy data.

As $t \rightarrow \infty$, the probability of an incorrect decision

$$\begin{aligned} P\{\text{classify as 1} \mid Y(0) = y(0), C(0) = 1, C(t) = 2\} &\rightarrow P\{(\theta_2 + \alpha W - \theta_1)^2 < (\alpha W)^2\} \\ &= P\{(W + \frac{\theta_2 - \theta_1}{\alpha})^2 < W^2\} = P\{W > \frac{|\theta_2 - \theta_1|}{2\alpha}\}. \end{aligned}$$

Suppose now the Cauchy classification procedure is used on the Cauchy data with Item 2 associated with the Signature at time t and Item 1 associated with the Signature at time 0. The probability of an incorrect classification is $P\{\text{classify as 1} \mid Y(0) = y(0), C(0) = 1, C(t) = 2\}$

$$\begin{aligned} &= P\{\alpha(1 - |\rho|^t)[\alpha^2(1 - |\rho|^t)^2 + [\theta_2 + \alpha W - (\theta_1 + \rho^t(y(0) - \theta_1))]^2]^{-1} \\ &\quad > \alpha[\alpha^2 + (\theta_2 + \alpha W - \theta_1)^2]^{-1}\} \\ &= P\{(1 - |\rho|^t)[\alpha^2 + (\alpha W)^2] \\ &\quad > [\alpha^2(1 - |\rho|^t)^2 + [\theta_2 + \alpha W - (\theta_1 + \rho^t(y(0) - \theta_1))]^2]\}. \end{aligned} \tag{4.27}$$

As $t \rightarrow 0$, the probability of an incorrect classification

$$P\{\text{classify as 1} \mid Y(0) = y(0), C(0) = 1, C(t) = 2\} \rightarrow P\{0 > (\theta_2 + \alpha W - y(0))^2\} = 0.$$

Thus, the probability of an incorrect identification using the Cauchy procedure tends to 0 as $t \rightarrow 0$ for the case in which the Item associated with the

Signature at time t is different from the Item associated with the Signature at time 0.

As $t \rightarrow \infty$, the probability of an incorrect identification

$$P\{\text{classify as 1} \mid Y(0) = y(0), C(0) = 1, C(t) = 2\}$$

$$\rightarrow P\{W^2 > (W + \frac{(\theta_2 - \theta_1)}{\alpha})^2\} = P\{W > \frac{|\theta_2 - \theta_1|}{2\alpha}\}$$

the same as for the Gaussian procedure.

Hence, for the two limiting cases $t \rightarrow 0$ and $t \rightarrow \infty$ both the Cauchy and Gaussian procedures have the same misclassification probabilities for the case in which the Item associated with the Signature at time t is different than the one associated with the Signature at time 0. Note these are theoretical limiting results with all parameters known.

To further explore the differences between the Gaussian and Cauchy procedures for the scenario of different Items associated with Signatures and Cauchy data, let

$$g_t(W, y(0)) = [\alpha W + \theta_2 - \theta_1 - \rho^t(y(0) - \theta_1)]^2. \quad (4.28)$$

From (4.26) for the Gaussian procedure, the probability of an incorrect classification

$$P\{\text{classify as 1} \mid Y(0) = y(0), C(0) = 1, C(t) = 2\}$$

$$= P\{\sigma(\infty)^2 (1 - \rho^{2t}) \ln(1 - \rho^{2t}) < (\alpha W)^2 (1 - \rho^{2t}) - g_t(W, y(0))\}.$$

For the Cauchy procedure, the probability of an incorrect classification

$$P\{\text{classify as 1} \mid Y(0) = y(0), C(0) = 1, C(t) = 2\}$$

$$= P\{\alpha^2(1 - |\rho|^t)[1 - |\rho|^{t-1}] < (1 - |\rho|^t) (\alpha W)^2 - g_t(W, y(0))\}$$

$$= P\{\alpha^2(1 - |\rho|^t) |\rho|^t < (1 - |\rho|^t) (\alpha W)^2 - g_t(W, y(0))\}$$

$$\leq P\{\sigma(\infty)^2 (1 - \rho^{2t}) \ln(1 - \rho^{2t}) < (1 - |\rho|^t) (\alpha W)^2 - g_t(W, y(0))\}$$

$$\leq P\{\sigma(\infty)^2 (1 - \rho^{2t}) \ln(1 - \rho^{2t}) < (1 - \rho^{2t}) (\alpha W)^2 - g_t(W, y(0))\}$$

for t sufficiently small. Thus, for small t the Gaussian procedure will tend to have more incorrect classifications than the Cauchy procedure for the scenario of Cauchy data with the Item associated with the Signature at time t being different than the one associated with the signature at time 0. This effect is made stronger by the fact that if the Gaussian procedure is used then an estimate of $\sigma(\infty)^2$ will be needed. An estimate of $\sigma(\infty)^2$ for Cauchy data will tend to be very large since the Cauchy distribution does not have a finite variance. This effect will be seen in the simulations of the next subsection.

4.3 Results of simulation experiments

This subsection reports on results of simulation experiments to assess the behavior of the Gaussian and Cauchy classification procedures when they are confronted with data from the other distribution. For simplicity we assume there are two Items. In the first subsection the autoregressive process producing the data is Gaussian. In the second subsection the autoregressive process producing the data is Cauchy. In both subsections classification procedures using both the Cauchy and Gaussian distributional assumptions are assessed. In all cases $\rho_1 = \rho_2 = 0.5$, $\theta_1 = 1$, $\theta_2 = 2$. The simulations use the LLRANDOM random number generator; cf. Lewis and Uribe [1981].

a. Gaussian Data

The simulation in this subsection uses data from a Gaussian autoregressive process. We will assume the means of the two Signatures, θ_1 and θ_2 , are known and $\rho_1 = \rho_2 = \rho$ is also known. It remains to assess values for the (presumed known) scale parameters of the two classification procedures. In particular what should the scale parameter $\alpha = \alpha_1 = \alpha_2$ of the Cauchy procedure be when it is applied to Gaussian data? To obtain reasonable values for $\sigma_1 = \sigma_2 = \sigma$ for the Gaussian classification procedure and $\alpha = \alpha_1 = \alpha_2$ for the Cauchy classification procedure, the following simulation experiment was performed. The experiment has 100 replications. In each replication 100 independent, standard normals are generated. For each replication, the standard deviation of the data is computed and the maximum likelihood estimate of α is obtained numerically assuming a Cauchy density function of the form

$$f(x) = \frac{1}{\pi} \frac{\alpha}{\alpha^2 + x^2}, \quad -\infty < x < \infty.$$

The medians of the 100 estimates of α and the 100 standard deviations are calculated. The values obtained are $\hat{\sigma}_M = 1.0$ $\hat{\alpha}_M = 0.607$. Note that the estimates of α are using the incorrect model assumption of Cauchy for the Gaussian data. The value of $\hat{\sigma}_M$ is used in the Gaussian procedure to classify observations. The value of $\hat{\alpha}_M$ is used in the Cauchy procedure.

Tables 1 and 2 show results for simulation experiments with 500 replications. In each replication $Y(0)$ is generated from a normal distribution with mean θ_1 and standard deviation $\sigma(\infty) = \sigma/\sqrt{1-\rho^2}$ with $\sigma=1$ and $\rho = 0.5$. For Table 1 $Y(t)$ is generated from a normal distribution with mean

$$m(t) = \theta_1 + \rho^t(Y(0) - \theta_1)$$

and standard deviation $\sigma(t) = \sigma(\infty)\sqrt{1-\rho^t}$; namely the Signature observed at time t is from Item 1. For Table 2 $Y(t)$ is generated from a normal distribution with mean θ_2 and standard deviation $\sigma(\infty)$; namely the Signature at time t is from Item 2.

In both Tables the Gaussian classification procedure assumes $\hat{\sigma}_M = 1.0$ is the correct standard deviation. The Cauchy classification procedure assumes $\hat{\alpha}_M = 0.607$ is the correct value for α .

The values in Table 1 suggest that when the same Item is producing the Signature at time 0 and t , then the Gaussian procedure produces more correct classifications for small time t . However, the number of correct classifications is the same for both procedures for larger t .

The values of Table 2 suggest that if a different Item is producing the Signature at time t , then the Cauchy classification procedure has more correct classifications at time t for small t even though the data are Gaussian. For larger t , both procedures have the same number of correct identifications.

b. Cauchy Data

In this subsection the data arise from a Cauchy autoregressive process. The mean Signatures of the two Items, θ_1 and θ_2 , are assumed known and $\rho = \rho_1 = \rho_2$ is also assumed known. It remains to assess values for the scale parameters of the Gaussian and Cauchy classification procedures. In particular, what should the scale parameter $\sigma = \sigma_1 = \sigma_2$ of the Gaussian procedure be when it is applied to Cauchy data?

TABLE 1. GAUSSIAN DATA

Item 1 Produces the Signature at time 0

Item 1 Produces the Signature at time t

Time t	Fraction Correct Identifications		Number of times	
	Gaussian Proc.	Cauchy Proc.	Gaussian Correct /Cauchy Incorrect	Gaussian Incorrect/Cauchy Correct
1	0.77	0.65	50	0
2	0.68	0.67	5	0
5	0.67	0.67	0	0
10	0.70	0.70	0	0

TABLE 2. GAUSSIAN DATA

Item 1 Produces the Signature at time 0

Item 2 Produces the Signature at time t

Time	Fraction Correct Identifications		Number of times	
	Gaussian Proc.	Cauchy Proc.	Gaussian Correct /Cauchy Incorrect	Gaussian Incorrect/Cauchy Correct
1	0.64	0.71	0	38
2	0.69	0.71	0	6
5	0.64	0.64	1	0
10	0.68	0.68	0	0

To obtain reasonable values for σ for the Gaussian classification procedure and α for the Cauchy classification procedure, the following simulation experiment was performed. The experiment has 100 replications. Each replication generates 100 standard Cauchy random numbers. For each replication the standard deviation of the data is computed and the maximum likelihood estimate of α is obtained numerically. The medians of the 100

estimates of α and the 100 standard deviations are computed. The values obtained are

$$\hat{\sigma}_M = 13.23 \text{ and } \hat{\alpha}_M = 1.03.$$

Note the high value of the standard deviation.

Tables 3 and 4 present results of simulation experiments in which the data are from a Cauchy autoregressive process. All experiments have 500 independent replications. For each replication $Y(0)$ is generated from a Cauchy distribution with location parameter θ_1 , and scale parameter 1; that is, Item 1 is producing the Signature at time 0. For replications reported in Table 4, $Y(t)$ is generated from a Cauchy distribution with location parameter θ_2 and scale parameter 1; that is, Item 2 is producing the Signature at time t . For replications reported in Table 3, $Y(t)$ is generated from a Cauchy distribution having density function

$$f(x) = \frac{1}{\pi} \frac{\alpha(t)}{\alpha(t)^2 + (x - m(t))^2}$$

with

$$m(t) = \theta_1 + \rho^t(y(0) - \theta_1)$$

and

$$\alpha(t) = (1 - |\rho|^t);$$

that is, Item 1 is also producing the Signature at time t .

In both Tables 3 and 4, the Gaussian classification procedure assumes a standard deviations $\sigma_1 = \sigma_2 = \hat{\sigma}_M$. The Cauchy classification procedure assumes the α -parameters $\alpha_1 = \alpha_2 = \hat{\alpha}_M$.

The results of Table 3 indicate that for small times t , if the same Item is producing the Signature at time 0 and time t , then the Gaussian classification

procedure has more correct classifications even though the data are Cauchy. For larger times t , the number of correct classifications is the same for both procedures. On the other hand, the results of Table 4 indicate that if a different item is producing the Signature at time t , then the Cauchy classification procedure has many more correct classifications than the Gaussian procedure for small times t . Once again the number of correct identifications is the same for both procedures as t becomes larger.

TABLE 3. CAUCHY DATA

Item 1 Produces the Signature at time 0

Item 1 Produces the Signature at time t

Time	Fraction Correct Identifications		Number of times	
	Gaussian Proc.	Cauchy Proc.	Gaussian Correct /Cauchy Incorrect	Gaussian Incorrect/Cauchy Correct
1	0.98	0.73	124	0
5	0.72	0.69	18	0
10	0.66	0.66	0	1

TABLE 4. CAUCHY DATA

Item 1 Produces the Signature at time 0

Item 2 Produces the Signature at time t

Time	Fraction Correct Identifications		Number of times	
	Normal Proc.	Cauchy Proc.	Normal Correct /Cauchy Incorrect	Normal Incorrect/Cauchy Correct
1	0.09	0.74	0	316
2	0.14	0.70	0	282
5	0.67	0.61	0	29
10	0.64	0.64	0	0

c. Summary

The differences in performance of the two classification procedures appear for small time t . If the same Item is producing Signatures at both 0 and t , then the Gaussian classification procedure has more correct classifications for small times t for both Gaussian and Cauchy data. If a different Item is producing the Signature at time t , then the Cauchy classification procedure has more correct classifications for both Gaussian and Cauchy data. The effect is strongest if the data are from a Cauchy autoregressive process; in this case the Gaussian procedure does very poorly when different Items are producing the Signatures.

In summary, it is important to realize that the performance of a Bayesian classification procedure can be influenced by its underlying distributional assumptions. A classification procedure based on Gaussian distributional assumptions can be reluctant to classify a new observation coming from a different item as being associated with a new item. A classification procedure based on Cauchy distributional assumptions can be reluctant to classify a new observation which comes from the same item as that being associated with the same item. Hence, if there is uncertainty about the underlying distribution of the data, it might be better to combine results of several classification procedures based on different distributional assumptions.

REFERENCES

Donald P. Gaver and Patricia A. Jacobs. "Item identities and their related observables." Naval Postgraduate School Technical Report, NPS55-90-03, 1989.

P. A. W. Lewis and L. Uribe. "The new Naval Postgraduate School random number package LLRANDOM II." Naval Postgraduate School Technical Report, NPS-55-81-005, 1981.




Figure 1

INITIAL DISTRIBUTION LIST

1.	Library (Code 0142).....	2
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
2.	Defense Technical Information Center.....	2
	Cameron Station	
	Alexandria, VA 22314	
3.	Office of Research Administration (Code 012).....	1
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
4.	Prof. Peter Purdue.....	1
	Code OR-Pd	
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
5.	Department of Operations Research (Code 55).....	1
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
6.	Prof. Donald Gaver, Code OR-Gv.....	15
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
7.	Prof. Patricia Jacobs	15
	Code OR/Jc	
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
8.	Center for Naval Analyses.....	1
	4401 Ford Avenue	
	Alexandria, VA 22302-0268	
9.	Dr. David Brillinger.....	1
	Statistics Department	
	University of California	
	Berkeley, CA 94720	

10. Dr. F. Gnanadesikan1
 Bellcore
 435 South Street
 Morris Township NJ 07960

11. Prof. Bernard Harris1
 Dept. of Statistics
 University of Wisconsin
 610 Walnut Street
 Madison, WI 53706

12. Prof. I. R. Savage.....1
 Dept. of Statistics
 Yale University
 New Haven, CT 06520

13. Prof. W. R. Schucany.....1
 Dept. of Statistics
 Southern Methodist University
 Dallas, TX 75222

14. Prof. D. C. Siegmund1
 Dept. of Statistics
 Sequoia Hall
 Stanford University
 Stanford, CA 94305

15. Prof. H. Solomon1
 Department of Statistics
 Sequoia Hall
 Stanford University
 Stanford, CA 94305

16. Dr. Ed Wegman.....1
 George Mason University
 Fairfax, VA 22030

17. Dr. P. Welch1
 IBM Research Laboratory
 Yorktown Heights, NY 10598

18. Dr. Neil Gerr1
 Office of Naval Research
 Arlington, VA 22217

19. Prof. Roy Welsch.....1
Sloan School
M.I.T.
Cambridge, MA 02139
20. Dr. J. Abrahams, Code 1111, Room 6071
Mathematical Sciences Division, Office of Naval Research
800 North Quincy Street
Arlington, VA 22217-5000
21. Prof. J. R. Thompson.....1
Dept. of Mathematical Science
Rice University
Houston, TX 77001
22. Dr. P. Heidelberger1
IBM Research Laboratory
Yorktown Heights
New York, NY 10598
23. Prof. M. Leadbetter...1
Department of Statistics
University of North Carolina
Chapel Hill, NC 27514
24. Prof. D. L. Iglehart1
Dept. of Operations Research
Stanford University
Stanford, CA 94350
25. Prof. J. B. Kadane.....1
Dept. of Statistics
Carnegie-Mellon University
Pittsburgh, PA 15213
26. Prof. J. Lehoczky1
Department of Statistics
Carnegie-Mellon University
Pittsburgh, PA 15213
27. Dr. J. Maar (R513)1
National Security Agency
Fort Meade, MD 20755-6000

28. Prof. M. Mazumdar1
 Dept. of Industrial Engineering
 University of Pittsburgh
 Pittsburgh, PA 15235

29. Prof. M. Rosenblatt1
 Department of Mathematics
 University of California, San Diego
 La Jolla, CA 92093

30. Prof. H. Chernoff1
 Department of Statistics
 Harvard University
 1 Oxford Street
 Cambridge, MA 02138

31. Dr. T. J. Ott1
 Bellcore
 435 South Street
 Morris Township, NJ 07960

32. Dr. Alan Weiss1
 AT&T Bell Laboratories
 Mountain Avenue
 Murray Hill, NJ 07974

33. Prof. Joseph R. Gani1
 Mathematics Department
 University of California
 Santa Barbara, CA 93106

34. Prof. Frank Samaniego1
 Statistics Department
 University of California
 Davis, CA 95616

35. Dr. James McKenna1
 AT&T Bell Laboratories
 Mountain Avenue
 Murray Hill, NJ 07974

36. Commander.....1
Attn: G. F. Kramer
Naval Ocean Systems Center
Code 421
San Diego, CA 92152-5000

37. Prof. Tom A. Louis1
School of Public Health
University of Minnesota
Mayo Bldg. A460
Minneapolis, MN 55455

38. Dr. Nan Laird.....1
Biostatistics Dept.
Harvard School of Public Health
677 Huntington Ave.
Boston, MA 02115

39. Dr. Marvin Zelen.....1
Biostatistics Department
Harvard School of Public Health
677 Huntington Ave.
Boston, MA 02115

40. Dr. John Orav.....1
Biostatistics Department
Harvard School of Public Health
677 Huntington Ave.
Boston, MA 02115

41. Prof. R. Douglas Martin.....1
Department of Statistics, GN-22
University of Washington
Seattle, WA 98195

42. Prof. W. Stuetzle1
Department of Statistics
University of Washington
Seattle, WA 98195

43. Prof. F. W. Mosteller1
 Department of Statistics
 Harvard University
 1 Oxford St.
 Cambridge, MA 02138

44. Dr. D. C. Hoaglin1
 Department of Statistics
 Harvard University
 1 Oxford Street
 Cambridge, MA 02138

45. Prof. N. D. Singpurwalla1
 George Washington University
 Washington, DC 20052

46. Prof. George S. Fishman1
 Curr. in OR & Systems Analysis
 University of North Carolina
 Chapel Hill, NC 20742

47. Dr. Alan F. Petty1
 Code 7930
 Naval Research Laboratory
 Washington, DC 20375

48. Prof. Bradley Efron1
 Statistics Dept.
 Sequoia Hall
 Stanford University
 Stanford, CA 94305

49. Prof. Carl N. Morris1
 Statistics Department
 Harvard University
 1 Oxford St.
 Cambridge, MA 02138

50. Dr. John E. Rolph1
 RAND Corporation
 1700 Main St.
 Santa Monica, CA 90406

51. Prof. Linda V. Green1
Graduate School of Business
Columbia University
New York, NY 10027

52. Dr. David Burman.....1
AT&T Bell Telephone Laboratories
Mountain Avenue
Murray Hill, NJ 07974

53. Dr. Ed Coffman.....1
AT&T Bell Telephone Laboratories
Mountain Avenue
Murray Hill, NJ 07974

54. Prof. William Jewell.....1
Operations Research Department
University of California, Berkeley
Berkeley, CA 94720

55. Prof. D. C. Siegmund.....1
Dept. of Statistics, Sequoia Hall
Stanford University
Stanford, CA 94305

56. Operations Research Center, Rm E40-164.....1
Massachusetts Institute of Technology
Attn: R. C. Larson and J. F. Shapiro
Cambridge, MA 02139

57. Arthur P. Hurter, Jr.1
Professor and Chairman
Dept. of Industrial Engineering and Management Sciences
Northwestern University
Evanston, IL 60201-9990

58. Institute for Defense Analysis.....1
1800 North Beauregard
Alexandria, VA 22311

59. Prof. J. W. Tukey1
Statistics Dept., Fine Hall
Princeton University
Princeton, NJ 08540

60.	Dr. Daniel H. Wagner	1
	Station Square One	
	Paoli, PA 19301	
61.	Dr. Colin Mallows.....	1
	AT&T Bell Telephone Laboratories	
	Mountain Avenue	
	Murray Hill, NJ 07974	
62.	Dr. D. Pregibon.....	1
	AT&T Bell Telephone Laboratories	
	Mountain Avenue	
	Murray Hill, NJ 07974	
63.	Dr. Jon Kettenring.....	1
	Bellcore	
	435 South Street	
	Morris Township, NJ 07960	
64.	Prof. David L. Wallace	1
	Statistics Dept., University of Chicago	
	5734 S. University Ave.	
	Chicago, IL 60637	
65.	Dr. S. R. Dalal.....	1
	AT&T Bell Telephone Laboratories	
	Mountain Avenue	
	Murray Hill, NJ 07974	
66.	Dr. M. J. Fischer	1
	Defense Communications Agency	
	1860 Wiehle Avenue	
	Reston, VA 22070	
67.	Dr. Prabha Kumar	
	Defense Communications Agency.....	1
	1860 Wiehle Avenue	
	Reston, VA 22070	
68.	Dr. B. Doshi	1
	AT&T Bell Laboratories	
	HO 3M-335	
	Holmdel, NJ 07733	

69. Dr. D. M. Lucantoni.....1
AT&T Bell Laboratories
Holdmel, NJ 07733
70. Dr. V. Ramaswami.....1
MRE 2Q-358
Bell Communications Research, Inc.
435 South Street
Morristown, NJ 07960
71. Prof. G. Shantikumar.....1
The Management Science Group
School of Business Administration
University of California
Berkeley, CA 94720
72. Dr. D. F. Daley.....1
Statistic Dept. (I.A.S.)
Australian National University
Canberra, A.C.T. 2606
AUSTRALIA
73. Dr. Guy Fayolle.....1
I.N.R.I.A.
Dom de Voluceau-Rocquencourt
78150 Le Chesnay Cedex
FRANCE
74. Professor H. G. Daellenbach.....1
Department of Operations Research
University of Canterbury
Christchurch, NEW ZEALAND
75. Koh Peng Kong.....1
OA Branch, DSO
Ministry of Defense
Blk 29 Middlesex Road
SINGAPORE 1024
76. Professor Sir David Cox.....1
Nuffield College
Oxford, OXI INF
ENGLAND

77. Dr. A. J. Lawrence.....1
 Dept. of Math Stat., Univ. of Birmingham
 P. O. Box 363
 Birmingham B15 2TT
 ENGLAND
78. Dr. F. P. Kelly.....1
 Statistics Laboratory
 16 Mill Lane
 Cambridge
 ENGLAND
79. Dr. R. J. Gibbens.....1
 Statistics Laboratory
 16 Mill Lane
 Cambridge
 ENGLAND
80. Dr. John Copas.....1
 Dept. of Math Stat., Univ. of Birmingham
 P. O. Box 363
 Birmingham B15 2TT
 ENGLAND
81. Dr. D. Vere-Jones.....1
 Dept. of Math, Victoria Univ. of Wellington
 P. O. Box 196
 Wellington
 NEW ZEALAND
82. Prof. Guy Latouche1
 University Libre Bruxelles
 C. P. 212, Blvd. De Triomphe
 B-1050 Bruxelles
 BELGIUM
83. Naval Ocean Systems Center1
 San Diego, CA 92152

DUDLEY KNOX LIBRARY



3 2768 00337195 6